



Food puzzles for cats: feeding for physical and emotional wellbeing

Leticia MS Dantas¹, Mikel M Delgado^{2,3}, Ingrid Johnson⁴
 and CA Tony Buffington⁵

Journal of Feline Medicine and Surgery
 1–12

© The Author(s) 2016

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/1098612X16643753

jfms.com

This paper was handled and
 processed by the American Editorial
 Office (AAFP) for publication in JFMS



Abstract

This article aims to equip veterinary professionals with the tools to assist clients in the use of food puzzles for their cats as ways to support feline enrichment, physical health and emotional wellbeing. We outline the types of food puzzles, how to introduce them to cats and how to troubleshoot challenges with their use. Owing to the paucity of evidence-based studies of food puzzles, we provide examples of the use and benefits of food puzzles from our own veterinary and behavioral practices.

Accepted: 17 March 2016

Introduction

Although cats are currently the most commonly kept pet in the USA, the conditions they are kept in are perhaps the least natural to them, especially given that the domestic cat's behavior and behavioral needs are very similar to those of their closest wild ancestor, the African wildcat.¹ Current veterinary and cat care guidelines (eg, American Veterinary Medical Association; Indoor Pet Initiative; American Association of Feline Practitioners) encourage keeping cats indoors for safety, health and ecological reasons, but this recommendation, along with the concurrent misperception of cats as low-maintenance pets, means that many cats are housed in suboptimal environments. One significant influence on cats' living conditions is how they are routinely fed. Most cats are offered food ad libitum from a bowl, are often required to share feeding areas or dishes with other cats and have to expend little to no effort to acquire calories.

Cats are natural predators that tend to eat multiple small meals each day in nature.^{2,3} When able to hunt, cats make several hunting attempts each day, only approximately half of which lead to a prey item.⁴ Indoor housing has been associated with increases in the occurrence of obesity,^{5–7} type 2 diabetes mellitus,⁸ joint problems⁹ and chronic lower urinary tract signs.¹⁰ Furthermore, the risk of behavioral and mental health problems may increase with confinement.¹¹ Commonly reported behavioral concerns from cat owners include aggression, attention-seeking behaviors and stress-related behaviors such as house soiling and overgrooming.^{12,13} These behavior problems can lead to a weakening of the human–animal

bond, and in many cases result in unwarranted euthanasia of the cat.^{14,15} Environmental enrichment may have some mitigating effects on these stress-related behaviors.¹⁶

We propose that one approach to environmental enrichment is to take advantage of cats' natural instinct to work for their food. In this review, we will explain what food puzzles are and why they are a biologically relevant enrichment device for cats. We will provide an overview of the most common types of food puzzles, how they should be used and how to overcome any client resistance to changing the method of food delivery to their cats. The effect of food puzzles on cats is a relatively new area of study, so in addition to reviewing the existing empirical evidence, we will provide case studies from our professional practices showing health and behavioral benefits from their use. Finally, we provide tools for

¹Behavioral Medicine Service, University of Georgia Veterinary Teaching Hospital, Athens, GA, USA

²Department of Psychology, University of California, Berkeley, CA, USA

³Feline Minds, Richmond, CA, USA

⁴Fundamentally Feline, Atlanta, GA, USA

⁵Veterinary Clinical Sciences, The Ohio State University, Columbus, OH, USA

Corresponding author:

Mikel M Delgado BA, Department of Psychology, University of California, Mailcode 1650, Berkeley, CA 94720-1650, USA
 Email: mikeldelgado@berkeley.edu

assessing the most appropriate types of food puzzles for individual cats, and a user-friendly handout (Supplementary material) that practitioners can give to their clients to get them started.

What are food puzzles?

Food puzzles were originally created to provide enrichment for captive zoo and laboratory animals.¹⁷ They typically consist of any object that can hold food and be manipulated to release food when the animal interacts with it. Food puzzles may be mobile (rolled or pushed) or stationary, and they can be used to provide either wet or dry food. They may be purchased or homemade (see Figure 1).

Mobile food puzzles are often shaped like a ball, egg or tube. Their rounded surfaces make the puzzles easy for cats to roll by pushing with a paw or their nose. These puzzles typically have one or more holes in them, and in some cases these puzzles can be adjusted to make release of food easier or harder by changing the size or number of open spaces that can dispense food. The current

iterations of these puzzles are typically designed for use with dry food or treats.

Stationary puzzles are typically larger, with sturdy bases, and holes, cups or channels. Dry food can be placed in the holes and cups, which must be fished out with a paw. Wells may be filled with wet food; the cat must lick food out of these wells, mimicking how cats use their jaw muscles to remove flesh from bone.

Homemade puzzles for dry food can be made easily by cutting holes in containers such as yogurt containers, toilet paper rolls, egg cartons, margarine tubs or water bottles. Ice cube trays or muffin pans can be used for wet food, and yogurt lids can be placed over the individual reservoirs or cups to increase difficulty.

Benefits of using food puzzles

Zoos and sanctuaries encounter many obstacles to providing adequate housing for felid species, which may have difficulty adjusting to captivity for several reasons. Territory and hunting opportunities are restricted, and many solitary species are housed in pairs or in groups.¹⁸ The parallels with domestic cat housing are numerous.



Figure 1 Clockwise from upper-left hand corner: homemade mobile, homemade stationary, purchased stationary and purchased mobile food puzzles. *Photos courtesy of Ingrid Johnson, Leticia Dantas and Mikel Delgado*

Implementing enrichment by providing foraging opportunities and food puzzles offers several benefits to captive large cats, including reducing stereotypies such as pacing,^{19,20} improving body condition²¹ and increasing exploratory behavior.^{18,22}

Current guidelines for the care and welfare of domestic cats suggest that they be allowed to express the predatory sequence to the extent possible, including active acquisition of food.^{4,23,24} Although few empirical studies of the benefits of food puzzles for companion animals have been conducted, provision of food puzzles has been shown to increase activity and reduce problematic behavior in dogs.²⁵ In cats, various forms of enrichment (such as play, perches, play towers and novel toys) have been shown to reduce signs of stress¹⁶ and to contribute to weight loss.²⁶ Table 1 outlines cases from our veterinary and behavioral practices that resulted in either behavioral or health-related benefits after the implementation of food puzzles along with other forms of behavior modification. Benefits we have observed include weight loss, decreased aggression toward humans and other cats, reduced anxiety and fear, cessation of attention-seeking behaviors and resolution of litter box avoidance.

Furthermore, implementing food puzzles provides few risks of decreasing the welfare of cats. In one study, 85% of cats (23/27) engaged with a food puzzle, without increases in aggression between group-housed cats in a shelter,²⁷ suggesting few problems with implementing food puzzles in multi-cat environments. Although problem-solving to acquire food may initially frustrate some animals, presenting animals with some level of challenge that is appropriate to their natural ecology and matched to their skill level is likely to provide cognitive, physical and behavioral benefits in otherwise-enriched surroundings.²⁸

Implementing food puzzles

As with the introduction of any new resource, successfully implementing food puzzles requires some planning. Clients should be prepared to try a few different types of food puzzles, because cats may have individual preferences for the type of puzzle or how they interact with puzzles (eg, some cats prefer mobile puzzles that can be pushed or rolled, while others are more adept at stationary puzzles; some cats are more likely to use their paws to move a toy, while others may push the toy with their nose). In Table 2, we have provided a guide to helping clients choose a starter puzzle. Furthermore, because implementing food puzzles offers enrichment beyond just as a means of providing food, the end goal is to have several different types of puzzles available for cats (as is recommended with other toys). The most important key to success is for clients to introduce puzzles to the cats correctly. This means setting the difficulty level to meet the abilities of the cat, and by increasing the cat's motivation to interact with

the puzzle as much as possible. Table 3 includes a summary of how best to introduce food puzzles.

Initially, obtaining food from the puzzle needs to be as easy as obtaining food from the food bowl. This means that the cat should have to do very little work for food at first. The puzzle should be filled as much as possible, and should have several, large holes to allow food to fall out easily. The puzzle should roll with little manipulation. For stationary puzzles, cups or reservoirs should be overflowing (see Figure 2).

As cats may initially be resistant to working for food, particularly if they have a history of not having to do so,^{29,30} the regular food dish may need to be removed when introducing food puzzles. Because some cats may view their food container as a 'safety signal', or may be stressed by changes in their environment, some cats prefer having the puzzle initially introduced as a choice next to the usual feeding container at the time of feeding, and containing the cat's usual food. Some cats, particularly those that tend to be nervous, may prefer that the food puzzle be placed in a quiet area, where they can explore it undisturbed.

The food puzzle should be introduced when the cat is likely to be hungry. Motivation may be increased at first by using a novel food type in the puzzle, such as treats or a dental diet. As the cats become more adept at using the puzzle, the food can be changed to their regular diet, or a mix of their regular dry food and treats.

For dry food puzzles, the client should place food on the floor next to and around the puzzle and allow the cat to eat around the puzzle. Their cat may inadvertently move the puzzle while eating, which will help them make the association between moving the puzzle and receiving food. The owner may even gently roll or nudge the puzzle at first to maintain the cat's interest. Eventually, regular food dishes can be removed and the cat can receive all of its daily food from puzzles.

Troubleshooting potential challenges and solutions

Some clients (and cats) may be reluctant to accept introduction of food puzzles. Making food puzzles as user-friendly and convenient as possible increases the client's likelihood of using them, and, consequently, improves their cat's welfare. Our collective experience is that most, if not all, cats can adjust to food puzzles, given time, patience and proper staging of difficulty. Some common challenges to food puzzle use and how we address them in our veterinary and behavioral practices include the following.

The client does not think their cat will use food puzzles

We have not encountered cats that could not adapt to food puzzles. Senior cats, kittens, three-legged cats, blind cats and cats with other disabilities, such as partial paralysis, have all been observed to use a food puzzle of

Table 1 Case examples where food puzzles were implemented to aid with a health or behavioral concern

Breed	Sex	Age (years)	Presenting concern	Food puzzle(s) implemented	Other modifications implemented	Outcome
DMH	2 NM	11 and 6	Obesity	Rolling and static (purchased)	–	Older cat lost 6.4% of body weight in 3.5 months, and increased mobility Younger cat lost 11% of body weight in 12 months Weight loss: 11% in 12 months
DMH	SF	9	Obesity	Rolling (homemade and purchased)	–	Weight loss: 20% in 12 months
DSH	NM	8	Obesity	Rolling (homemade and purchased)	–	Cats were able to play with food puzzles simultaneously; older cat preferred static puzzles and younger cat preferred rolling puzzles. The altercations between cats decreased significantly
DSH	2 NM	7 and 1	Younger cat trying to play rough with older cat	Rolling and static (purchased)	A general enrichment plan and clicker training were also introduced	Food puzzles slowed down both cats' eating and a decrease in meowing for food was observed. Behavior around owner's preparation of food and while eating was improved as long as cats were distracted with a food puzzle.
DMH	NM	3.5	Sibling cats meowing for food, waking owner up to be fed, interfering with owner's preparation and eating of meals, stealing food from plates, sinks and counters	Rolling and static (homemade and purchased)	The owner was instructed to avoid leaving food and dirty dishes on counters and sinks to prevent cats from being inadvertently rewarded for undesirable behaviors	The use of puzzles immediately resolved the situations where the cat experienced frustration (anticipation of meals, attention-seeking) and the cat started to show appeasement behaviors as opposed to impulsively attacking. The aggressive behavior resolved within 6 months
British Shorthair	NM	3	Impulsive and frustration-based aggression towards owner (biting with no warning when anticipating meals, and when attention was not given)	Rolling (purchased) and static (homemade)	–	Introduction of puzzles associated with marked decrease in disorientation, improved sleep–wake cycle alterations and cessation of nocturnal vocalizations, as well as increased social behavior between the cat and other cats in the household (three NM and one SF in total) and the clients. Provision of enrichment avoided introduction of pharmacological treatment
DSH	NM	10	Disorientation, nocturnal vocalization, alterations in the sleep–wake cycle, decreased social interactions, diagnosed with cognitive dysfunction syndrome	Rolling and static (homemade and purchased)	–	The cat's fear-related aggressive behaviors gradually decreased over time in frequency and intensity. After 1 year of treatment, the client was able to pet and pick up the cat, and the cat came when called. No more episodes of aggression were reported
DSH	SF	8	Fear-based aggression toward owner (cat signaled fear when approached, showed severe avoidance behavior and bit if touched or picked up)	Rolling and static (purchased)	In addition to the puzzles provided, the client was instructed to desensitize and counter-condition the cat to the owner's approach, and to positively reinforce alternative behaviors	

(Continued)

Table 1 (Continued)

Breed	Sex	Age (years)	Presenting concern	Food puzzle(s) implemented	Other modifications implemented	Outcome
DSH	NM	9	Noise phobia (cat had fear and panic reactions to several types of sudden and loud noises, sometimes redirecting aggression to a nearby cat in the multi-cat household)	Rolling and static (purchased)	–	Redirected aggression resolved, signs of noise phobia significantly improved
DSH	NM	2	Fear of people (familiar and unfamiliar)	Rolling and static (purchased)	–	Behavior significantly improved, with cat showing attachment signs to both owners, coming when called, allowing (and being relaxed during) petting; avoidance behavior ceased
DSH	1 SF and 1 NM	8 and 2	Fear-based aggressive behavior (older toward younger cat); younger cat redirected aggression toward the older cat. Altercations frequently led to bites	Rolling and static (purchased)	–	Intensity and frequency of aggression decreased and no more bites occurred. Both cats started to use avoidance behavior rather than aggression. The younger cat stopped redirecting aggression toward the older cat
DSH	2 SF	10 and 8	Fear-based aggressive behavior toward cat housemate (younger cat would signal fear and attack the older cat whenever it approached)	Rolling and static (purchased). Puzzles were provided for both cats, which were temporarily separated	Desensitization and counter-conditioning therapy was implemented with the use of the rolling puzzles and during feeding sessions of highly palatable food	Fear signaling (hissing, avoidance) was still seen sporadically, but no episodes of offensive aggression were reported after 6 months of treatment
DSH	NM	7 months	Pouncing and stalking the clients' guinea pigs	Rolling and static (homemade)	Counter-conditioning therapy sessions toward the guinea pigs	Implementation of puzzles and desensitization and counterconditioning therapy sessions toward the guinea pigs resulted in cessation of the cat's behaviors toward them
DLH	NM	1	Anxiety signs when left alone (increased vocalizations, agitation, pacing), hyper-attachment (followed owner constantly, stress signs when owner was out of sight) and anxiety response (agitation and tension) to owner departure cue. Diagnosed with separation anxiety syndrome	Rolling and static (homemade and purchased)	A comprehensive enrichment plan was formulated for this cat, with the addition of a more complex environment (vertical and hiding areas)	The anxiety signs while the client was away completely stopped. Signs of hyper-attachment and anxiety responses to owner departure cues were still seen but gradually weaned within 1 year of treatment

(Continued)

Table 1 (Continued)

Breed	Sex	Age (years)	Presenting concern	Food puzzle(s) implemented	Other modifications implemented	Outcome
DLH	SF	9	Signs of depression when left alone by the main caregiver (anorexia, social withdrawal, lack of play behavior), hyper-attachment (following primary caregiver constantly, stress signs when owner was out of sight) and anxiety response to owner departure cue (agitation and tension), diagnosed with separation anxiety syndrome	Rolling and static (homemade and purchased)	A comprehensive enrichment plan was formulated for this cat, with the addition of a more complex environment (vertical and hiding areas). A safe place was also conditioned	The signs of depression while the primary caregiver was away and of hyper-attachment and anxiety responses to owner departure cues decreased in frequency with gradual improvement during 6 months. Within 1 year, the client ranked the cat's improvement as excellent (signs were mild or not seen)
Varied	2 SF, 3 NM	Varied	Multi-cat household, meal-related fighting and urine marking. Urine marking continued, despite the cats living in separate spaces, owing to fighting. Regurgitation of undigested food was observed during meals	Rolling and static (homemade and purchased)	Cats were gradually reintroduced using positive reinforcement and counter-conditioning techniques. A litter box cafeteria was offered, allowing the cats to choose their preference. Post-cafeteria litter boxes were added in previously soiled areas of the home. Scratching posts were placed in areas where litter boxes were not an option, providing alternative marking opportunities. Meal feeding ceased and canned food was offered in excess in multiple feeding stations. Interactive play at a minimum of once a day was recommended and implemented	Significant decrease in inter-cat aggression, urine marking ceased, cats were fully integrated and no longer required separation. Regurgitation due to overeating much reduced in frequency
DSH	SF	6 months	Urination outside of the litter box, urinating in bathtub daily	Rolling (purchased). Owners began leaving food puzzles in the bathtub for the cat to play with	Owners made simultaneous adjustments to the litter box that increased compliance	The cat stopped urinating outside of the litter box
Persian	SF	6	Urination outside of litter box, diagnosed with feline idiopathic cystitis or Pandora syndrome	Rolling and static (purchased)	After initial buprenorphine treatment for the acute presentation, a comprehensive plan of multimodal environmental modification was provided	The cat has been in remission for 2.5 years

(Continued)

Table 1 (Continued)

Breed	Sex	Age (years)	Presenting concern	Food puzzle(s) implemented	Other modifications implemented	Outcome
DSH	NM	11	Urine marking (cat lived in a multi-cat household and had episodes throughout its adult life)	Rolling and static (purchased)	Vertical and hiding spaces were also added to the house	Marking behavior ceased
DSH	SF	9	Urination outside of litter box due to chronic feline idiopathic cystitis, possible location and substrate preference for toileting behavior and litter box aversion	Rolling and static (purchased)	In addition to the food puzzles, appropriate litter box management was recommended	Behavior completely resolved within 3 months of treatment
DLH	NM	12	Urination outside of litter box due to chronic feline idiopathic cystitis	Rolling and static (purchased)	Besides the food puzzles, appropriate litter box management was recommended	Behavior completely resolved within 6 months of treatment
DSH	NM	2	Urination outside of litter box due to chronic feline idiopathic cystitis, possible location and substrate preference for toileting behavior and litter box aversion	Rolling and static (purchased)	Besides the food puzzles, appropriate litter box management was recommended	Behavior completely resolved within 3 months of treatment
Maine Coon	NM	8 months	Pronounced fear toward household dog (13-week-old Golden Retriever); mild inappropriate play behavior toward owners and stress-induced scratching	Rolling and static (purchased)	Additional recommendations: safe place conditioning and desensitization and counterconditioning to dog, plus additional physical enrichment in the house	All problem behaviors resolved
DMH	SF	16	Obesity, lack of interaction with owners, secluded itself in one room. Fearful and antisocial with new kittens in home despite positive slow introduction	Started with static and advanced to all types	Many other accommodations had already been implemented for this cat: vertical space, hiding places, heating pads, canned and dry food, attempts to engage in interactive play and supportive therapy for arthritis, with little to no change	Weight loss: 32% in 18 months. After implementing static food puzzles the cat started to lose weight and play with the owners more. It eventually became interested in the other foraging toys outside of 'its room' that the kittens were playing with. It eventually rejoined the household, using both static and stationary toys. It also will now share and forage side by side with the young cats

DMH = domestic mediumhair; NM = neutered male; SF = spayed female; DSH = domestic shorthair; DLH = domestic longhair

Table 2 Checklist for helping clients choose a starter puzzle*

Does the cat eat wet and dry food?	Wet food is best used with stationary puzzles. Dry food can be easily used with stationary or mobile puzzles
Is owner okay with food around the house?	Mobile puzzles may lead to more food scatter
Is the owner concerned about night-time noise/activity?	Stationary puzzles and puzzles made of softer plastics make less noise
How does the cat typically interact with toys – with their paws, nose?	If the cat typically interacts with toys using their paws, they will likely do well with stationary or mobile puzzles. Cats who primarily use their nose will do best with mobile puzzles at first
Does the owner have financial constraints?	If yes, recommend homemade puzzles
Does the owner have time constraints?	If yes, recommend store-bought puzzles

*Ultimately, the goal is for clients to try several types of puzzles with their cats. This checklist can be used to help them choose a starter puzzle that is most likely to work with their cat

Table 3 Summary of how best to introduce food puzzles

When introducing food puzzles, clients should:
1 Continue to offer some food in the cat’s food bowl
2 Adjust the food delivery rate of the puzzle so that obtaining food is easy until the cat learns how to use the device
3 Monitor the cat to ensure that the experience is stimulating and positive
4 Be encouraged to convey feedback to your staff on the cat’s progress
5 Gradually increase the amount of food provided via the food puzzle once the cat is using the food puzzle successfully and no signs of stress or anxiety are observed



Figure 2 Puzzles should initially be filled as much as possible. *Photos courtesy of Mikel Delgado*

some type. Reminding the client of the cat's natural lifestyle as a hunter that works for food may be helpful. Sharing and discussing the food puzzle handout (Supplementary material) with the client, as well as demonstrating how the puzzles work, will help clients get started. Providing coaching and encouragement during the implementation process reduces the likelihood of the client concluding that their cat is unwilling or unable to work for their food.

The owner will not make or purchase food puzzles

The best way to address this is to have food puzzles available for sale in the veterinary office, or to provide referrals to local pet stores that sell food puzzles. There are also many good resources online for creating food puzzles out of recyclable materials (such as yogurt containers and plastic water bottles).

The owner does not want to prepare food puzzles daily

If the client is willing, they can acquire several food puzzles to rotate. All dry food puzzles can be prepared once a week and stored in airtight storage bins.³¹

The owner is concerned about noise/night-time activity

Stationary puzzles can be used at night, or mobile food puzzles can be confined to areas away from the bedroom.

The owner is resistant to having food scattered around the home

The cat will likely eat most of the food dispensed by the puzzle. However, food puzzles can be used in select rooms (eg, bathroom, office, kitchen), or in more restricted areas such as bathtubs, laundry baskets, under-the-bed storage containers or in the lids of large storage totes. The downside of restricting the area the puzzle is used in is that it makes food easier to obtain, and reduces the cat's movement and activity, and bathtubs and containers may provide other challenges for any cat with a mobility issue (such as older, arthritic cats).

The owner has tried a food puzzle and the cat would not use it

The client should be encouraged to try again, this time empowered with more detailed instructions and specific guidance. Staff should be trained to guide clients through learning challenges. Initial difficulty of use and increasing motivation are two of the most common barriers to cats' willingness to use food puzzles. For slow starters, placing handfuls of dry food in locations frequented by the cat (condos, window sills, beds) allows the cat the chance to discover food in novel places. These cats

generally can transition from this to static puzzles located in these areas.

All cats should be monitored closely at first when introducing a food puzzle, as missing several meals could be dangerous to vulnerable cats, particularly those that are overweight or obese. If the owner is concerned, they can offer the cat a portion of their daily food in a bowl, and the rest can be foraged until the cat has adapted to the use of the puzzle. Alternatively, the cat could be fed wet food from a dish, and dry food exclusively from puzzles. The ultimate goal should be all food from puzzles.

Multiple cats in the home

All cats in the home can be acclimated to food puzzles. Because cats may have individual preferences, several types of food puzzles can be distributed throughout the home, and each cat should be provided with their own puzzle. If one type of puzzle is more popular, then the owner should provide multiple puzzles of that type. This prevents cats from having to compete over food resources, and from being forced to eat in the same area, a behavior that is unnatural to solitary hunters. However, some cats are willing to use a food puzzle together (see Figure 3).²⁷

Presence of dogs in the home

For homes with dogs, puzzles can be used in restricted areas (see above), or baby gates can be used to keep dogs out of certain areas of the home where the cats' food puzzles are kept. The baby gates should be placed at a height where the cat can either jump over or crawl under, but the dog cannot. Dogs also can be taught a 'leave it' verbal cue, and be provided with their own foraging toys in a separate area of the house, or if possible, outdoors.

The cat appears frustrated by the food puzzle

Frustration can occur in animals when a previously obtainable resource changes or becomes inaccessible. Frustration may have adaptive properties, as it can lead to the persistence of efforts and emitting of novel behaviors to access a resource or solve a problem.^{28,32} While such behaviors would promote successful use of food puzzles, frustration may lead to development of fearful or aggressive behaviors in some animals, particularly when a problem is insoluble.³³ We have not encountered cats that developed such negative behaviors after the introduction of food puzzles, because difficulty was staged appropriately. We recognize that frustration is possible, particularly in unenriched environments; we recommend that food puzzles be introduced as part of a multi-modal enrichment plan,^{34,35} and that our implementation recommendations be followed carefully.



Figure 3 Some cats will use food puzzles together. *Photos courtesy of Ingrid Johnson and Leticia Dantas*



Figure 4 Placing a small puzzle inside a larger puzzle increases challenge for cats. *Photo courtesy of Ingrid Johnson*



Figure 5 A piñata-style puzzle that hangs. *Photo courtesy of Ingrid Johnson*

The cat becomes bored with the food puzzles

Adjusting the difficulty to the ability of the cat is important. Unchallenging tasks may lead to boredom or apathy.²⁸ When the cat becomes competent with a particular food puzzle, clients can maintain their cat's interest by:

- 1 Trying new types of food puzzles (including different designs or materials) or changing the location the puzzle is used in
- 2 Increasing the difficulty of the puzzle: owners can do this by changing the number and/or size of the holes, or by using objects that are opaque and give fewer visual cues. They also can adjust how mobile the puzzle is by using weighted puzzles, cubes or objects that have textured sides. Changing the substrate the puzzle is offered on (eg, carpet instead of hard flooring) can increase the challenge
- 3 Placing a small puzzle inside a larger puzzle (see Figure 4)

- 4 Placing 'dummy' objects inside the puzzle such as toy balls or mice that slow the release of food
- 5 Using piñata-style puzzles that dangle or hang and require the cats to reach up to manipulate them (see Figure 5). Clients should be advised that hanging toys can present a choking hazard and should only be used with supervision.

Conclusions

Food puzzles enable cat owners to provide mental stimulation and exercise, and increase the activity of their pet cats, while taking advantage of the domestic cat's natural inclination to work (hunt) for their food. Food puzzles are relatively easy to implement, and there are few risks associated with their use. In fact, there are likely many health and behavioral benefits from implementing food puzzles. We have provided some examples of the potential benefits of their use in concert with other behavior modifications in the retrospective case studies outlined in Table 1.

We acknowledge the limitations to the cases we provided; they are retrospective in nature, and, in some cases, food puzzles were not the only intervention, so one cannot be sure of the relative effectiveness of each component of the treatment. We included these cases acknowledging that this limitation is also often found in primary care medicine, and in hopes of encouraging further research to determine the most appropriate roles of food puzzles in promoting health and welfare for confined cats. Prospective trials are greatly needed to help determine the relative effectiveness of different approaches to environmental enrichment for confined cats. We also did not include all cases seen that involved recommendations for the use of food puzzles, although we attempted to provide a range of the types of cases that might be commonly encountered in primary care veterinary practice. Veterinary practices are often the main source of information for many pet owners, and we have provided suggestions, tools and user-friendly information to help veterinary professionals make the recommendation of food puzzles a standard practice.

Conflict of interest The authors declare no potential conflict of interest with respect to the research, authorship, and/or publication of this article.

Funding This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. MMD was supported by an NSF GRFP fellowship.

Supplementary material

Using food puzzles with your cat' handout.

References

- 1 Rodan I and Heath S. Feline behavioral health and welfare. St Louis, MO: Elsevier Health Sciences, 2015.
- 2 Beaver BV. Feline behavior. St Louis, MO: Elsevier Health Sciences, 2003.
- 3 National Research Council (US). Ad Hoc Committee on Dog and Cat Nutrition. Nutrient requirements of dogs and cats. Washington, DC: National Academies Press, 2006.
- 4 Rochlitz I. A review of the housing requirements of domestic cats (*Felis silvestris catus*) kept in the home. *Appl Anim Behav Sci* 2005; 93: 97–109.
- 5 Allan F, Pfeiffer D, Jones B, et al. A cross-sectional study of risk factors for obesity in cats in New Zealand. *Prev Vet Med* 2000; 46: 183–196.
- 6 Robertson I. The influence of diet and other factors on owner-perceived obesity in privately owned cats from metropolitan Perth, Western Australia. *Prev Vet Med* 1999; 40: 75–85.
- 7 Scarlett J, Donoghue S, Saidla J, et al. Overweight cats: prevalence and risk factors. *J Int Assoc Stud Obes* 1994; 18: S22–S28.
- 8 Slingerland LI, Fazilova VV, Plantinga EA, et al. Indoor confinement and physical inactivity rather than the proportion of dry food are risk factors in the development of feline type 2 diabetes mellitus. *Vet J* 2009; 179: 247–253.
- 9 Laflamme DP. Understanding and managing obesity in dogs and cats. *Vet Clin N Am Small Anim Pract* 2006; 36: 1283–1295.
- 10 Buffington CT. Idiopathic cystitis in domestic cats – beyond the lower urinary tract. *J Vet Intern Med* 2011; 25: 784–796.
- 11 Stella JL, Lord LK and Buffington CT. Sickness behaviors in response to unusual external events in healthy cats and cats with feline interstitial cystitis. *J Am Vet Med Assoc* 2011; 238: 67–73.
- 12 Porters N, De Rooster H, Verschueren K, et al. Development of behavior in adopted shelter kittens after gonadectomy performed at an early age or at a traditional age. *J Vet Behav* 2014; 9: 196–206.
- 13 Amat M, de la Torre JLR, Fatjó J, et al. Potential risk factors associated with feline behaviour problems. *Appl Anim Behav Sci* 2009; 121: 134–139.
- 14 Salman MD, New JG, Scarlett JM, et al. Human and animal factors related to relinquishment of dogs and cats in 12 selected animal shelters in the United States. *J Appl Anim Welfare Sci* 1998; 1: 207–226.
- 15 Salman MD, Hutchison J, Ruch-Gallie R, et al. Behavioral reasons for relinquishment of dogs and cats to 12 shelters. *J Appl Anim Welfare Sci* 2000; 3: 93–106.
- 16 Buffington CT, Westropp JL, Chew DJ, et al. Clinical evaluation of multimodal environmental modification (MEMO) in the management of cats with idiopathic cystitis. *J Feline Med Surg* 2006; 8: 261–268.
- 17 Young RJ. Environmental enrichment for captive animals. Hoboken, NJ: Blackwell Publishing, 2003.
- 18 Mellen JD and Shepherdson DJ. Environmental enrichment for felids: an integrated approach. *Int Zoo Yearbook* 1997; 35: 191–197.

- 19 Bashaw MJ, Blooms Smith MA, Marr M, et al. **To hunt or not to hunt? A feeding enrichment experiment with captive large felids.** *Zoo Biol* 2003; 22: 189–198.
- 20 Markowitz H and LaForse S. **Artificial prey as behavioral enrichment devices for felines.** *Appl Anim Behav Sci* 1987; 18: 31–43.
- 21 Young RJ. **The importance of food presentation for animal welfare and conservation.** *Proc Nutr Soc* 1997; 56: 1095–1104.
- 22 Williams B, Waran N, Carruthers J, et al. **The effect of a moving bait on the behaviour of captive cheetahs (*Acinonyx jubatus*).** *Anim Welfare* 1996; 5: 271–281.
- 23 Ellis SL. **Environmental enrichment: practical strategies for improving feline welfare.** *J Feline Med Surg* 2009; 11: 901–912.
- 24 Ellis SL, Rodan I, Carney HC, et al. **AAFP and ISFM feline environmental needs guidelines.** *J Feline Med Surg* 2013; 15: 219–230.
- 25 Schipper LL, Vinke CM, Schilder MBH, et al. **The effect of feeding enrichment toys on the behaviour of kennelled dogs (*Canis familiaris*).** *Appl Anim Behav Sci* 2008; 114: 182–195.
- 26 Clarke D, Wrigglesworth D, Holmes K, et al. **Using environmental and feeding enrichment to facilitate feline weight loss.** *J Anim Physiol Anim Nutr* 2005; 89: 427.
- 27 Dantas-Divers LM, Crowell-Davis SL, Alford K, et al. **Agonistic behavior and environmental enrichment of cats communally housed in a shelter.** *J Am Vet Med Assoc* 2011; 239: 796–802.
- 28 Meehan CL and Mench JA. **The challenge of challenge: can problem solving opportunities enhance animal welfare?** *Appl Anim Behav Sci* 2007; 102: 246–261.
- 29 Koffer K and Coulson G. **Feline indolence: cats prefer free to response-produced food.** *Psychon Sci* 1971; 24: 41–42.
- 30 Osborne SR. **The free food (contrafreeloading) phenomenon: a review and analysis.** *Anim Learn Behav* 1977; 5: 221–235.
- 31 Fundamentally Feline. **Implementing foraging as a feeding protocol.** <http://www.fundamentallyfeline.com/implementing-foraging-as-a-feeding-protocol> (2014, accessed January 31, 2016).
- 32 Wong PT **Frustration, exploration, and learning.** *Can Psychol Rev* 1979; 20: 133–144.
- 33 van Kampen HS. **Violated expectancies: cause and function of exploration, fear, and aggression.** *Behav Process* 2015; 117: 12–28.
- 34 Herron ME and Buffington CA. **Environmental enrichment for indoor cats: implementing enrichment.** *Compend Contin Educ Vet* 2012; 34: E1–E5.
- 35 Herron ME and Buffington CAT. **Environmental enrichment for indoor cats.** *Compend Contin Educ Pract Vet* 2010; 32: E1–E5.